

AP2005052124 13 JUL 2006

5

**MODULATION OF ENZYME ACTIVITY BY CHANGING
RECIPROCAL STEREO-GEOMETRIC POSITIONS OF TWO
CATALYTIC CENTERS IN AN ENZYME**

10

BACKGROUND

Recombination between sequences on two distinct duplex
DNAs, or between sequences, which are separated by an
15 intervening sequence on the same DNA duplex, occurs in all cells
capable of self-replication. Four-way Holliday junctions in DNA are a
feature of recombination reactions and are generated in both
homologous and site-specific recombination reactions (Lilley, D.M.J.
PNAS 94, 9513-9515 (1997)). The penultimate stage of
20 recombination involves resolution of the four-way junction by
catalysis. The catalysis is achieved by structure-specific nucleases
identified as resolvases (Aravind, L., et al. *Nuc. Acid Res.* 28:3417-
3432 (2000)). Resolvases are widespread in cells and additionally
are expressed by viruses. Resolvases have diverse properties. The
25 crystal structures of a number of resolvases have been reported,
including that of T7 Endonuclease I (T7 Endo I) (Hadden, J.M., et
al. *Nat. Struct. Biol.* 8:62-67 (2001)).

T7 Endo I has two catalytic centers that are juxtaposed in a
30 way that prevents the enzyme from forming a productive complex
with regular linear DNA but enables it to specifically bind and cleave
branched, perturbed or flexible DNA. The three-dimensional
structure of T7 Endo I shows that the two catalytic domains are well
separated and connected only by the β -sheet bridge (Hadden, J.M.,
35 et al. *Nat. Struct. Biol.* 8:62-67 (2001)). The bridge forms part of
an extended and tightly associated anti-parallel β -sheet (β 2).

5

T7 Endo I is a stable homodimer of 149 amino acid subunits (Parkinson, M.J. and Lilley, D.M.J. *J. Mol. Biol.* 270:169-178 (1997)). It is very basic (pI calc 9.5) and binds tightly (K_d 2nM) to four-way junctions in dimeric form. T7 Endo I resolves four-way
10 junctions by simultaneously introducing two nicks on the two continuous strands, at sites 5' to the junction (Déclais, A., et al. *EMBO J.* 22:1398-1409 (2003)). This structural analysis shows that T7 Endo I forms an intimately associated symmetrical homodimer comprising two catalytic domains connected by a bridge. Each
15 catalytic domain is composed of residues 17-44 from one subunit and residues 50-145 from the other.

Resolvases including T7 Endo I are structure-specific endonucleases capable of cleaving a broad range of DNA molecules
20 with a variety of structures such as branched structures and single-base mismatched heteroduplexes (Mashal, R.D., et al. *Nat. Genet.* 9:177-183 (1995)). This broad substrate specificity makes it difficult to identify the common structural features of the substrate that the enzyme selectively recognizes and cleaves (White, M.F., et
25 al. *J. Mol. Biol.* 269:647-664 (1997)). The broad substrate specificity may also contribute to the toxicity of the resolvase in host cells.

It would be desirable to reduce the toxicity of the enzymes having a similar structure to T7 Endo I to facilitate their
30 overproduction for use as reagents in molecular biology. It would also be desirable to selectively favor specific enzyme activities.

5

SUMMARY

An embodiment of the invention provides a modified DNA cleaving enzyme having at least 35% amino acid sequence identity with T7 Endo I, two catalytic centers separated by a β -bridge, and a mutation of at least one amino acid in the β -bridge that has an effect of altering cleavage activity.

In an embodiment of the invention, a feature of the modified DNA cleaving enzyme is reduced toxicity in a host cell permitting over-expression of the DNA cleaving enzyme.

In other embodiments of the invention, the modified DNA cleaving enzyme is capable of at least one of: cleaving cruciform structures in DNA; cleaving DNA at a site flanking a mismatch base pair; cleaving DNA in a non-sequence specific manner; nicking in a non-sequence dependent manner; and nicking opposite a pre-existing nick site. Cleavage on both strands of a DNA duplex results in a single stranded overhang of less than 11 nucleotides.

In comparison with the unmodified enzyme, the altered enzyme cleavage activity of the modified enzyme may, in certain embodiments, include a broadened enzyme specificity. For example, for the enzyme having a mutation in the β - bridge, DNA cleaving activity can occur at a site flanking a mismatch in a duplex after recognition of a mismatch containing any of A, T, G or C bases.

In another embodiment of the invention, an alteration in the enzyme cleavage activity of the modified enzyme compared to the unmodified enzyme occurs in a manganese-containing buffer. For

5 example, the alteration in cleavage activity may be one of:
maintenance of cleavage activity; reduction of non-specific nuclease
activity; enhanced nicking activity opposite a pre-existing nick site;
and a decreased ratio of nicking to double strand cleavage.

10 In another embodiment of the invention, an alteration in the
enzyme cleavage activity of the modified enzyme compared to the
unmodified enzyme occurs in a magnesium-containing buffer.
Accordingly, the observed altered enzyme activity is selected from:
an increased ratio of nicking of a cruciform structure in the DNA
15 relative to double strand cleavage; an increased ratio of cleaved
DNA of a cruciform to non-cleaved DNA; a reduced ratio of non-
specific nuclease activity; and a reduction in nicking opposite a pre-
existing nick site.

20 The altered enzyme activity of the modified enzyme can be
enhanced or reduced activity by modifying reaction conditions. For
example, modified reaction conditions may result from changing at
least one of: pH, temperature, addition of manganese ions to a
magnesium containing buffer or vice versa or changing the
25 concentration of magnesium or manganese ions in the reaction
mixture; and time of incubation of reactants.

In an embodiment of invention, the DNA cleaving enzyme is
selected from gene 3 (enterobacteriophage T7), T7
30 endodeoxyribonuclease I, Yersinia pestis phage phiA1122
endonuclease, Phage Phi Ye03-12 endonuclease, Phage T3
endonuclease, phage T3 endodeoxyribonuclease, Pseudomonas
phage gh-1 endonuclease, pseudomonas putida KT2440
endodeoxyribonuclease I; and Roseophage S101 RP endonuclease I.

5

In an embodiment of the invention, the modified enzyme has a mutation at a PA site in the β -bridge. For example, this mutation may be a deletion (Δ PA (where (Δ) corresponds to "deletion"), or a substitution of PA such as a single amino acid, a dipeptide, a tripeptide or a tetrapeptide substitution. Examples of substitutions include: PA/A, PA/AA, PA/PGA, PA/PAPA, PA/K, PA/G, PA/D and PA/P (where PA/A corresponds to a substitution(/) of PA dipeptide with A amino acid).

15 In an embodiment of the invention, the PA dipeptide is located at position 46 and 47 in SEQ ID. No. 12.

In an embodiment of the invention, a nucleic acid is provided that includes a DNA sequence that substantially corresponds to SEQ ID NO:1 wherein at least one mutation has been introduced into the nucleotide sequence corresponding to the β -bridge. For example, the mutation may be included at a site, which encodes the PA in the β -bridge. In this example, the mutation at the site that encodes PA may be any of a substitution or deletion such that the substituted nucleic acid encodes a single amino acid, a dipeptide, a tripeptide and a tetrapeptide. In a further example, the at least one mutation results in an amino acid change selected from PA/A, PA/AA, PA/PGA, PA/PAPA, Δ PA, PA/K, PA/G, PA/D and PA/P.

30 In embodiments of the invention, a vector is provided that includes any of the above examples of nucleic acids. In addition, a host cell is provided that includes any vector that includes a nucleic acid as described above.

5 In a further embodiment of the invention, a kit is provided that contains at least one of: a DNA cleaving enzymes in which a mutation has been introduced into the β -bridge as described above, a nucleic acid encoding the enzyme, a vector containing the nucleic acid, or a host cell containing the vector described above.

10

 In a further embodiment of the invention, a method is provided for modifying enzyme catalytic activity that includes: selecting an enzyme having two catalytic centers connected by a β -bridge, the catalytic centers being located at reciprocal stereo-
15 geometric positions in the enzyme; and changing the reciprocal stereo-geometric position of the two catalytic centers by introducing a mutation into the β -bridge.

 In a further embodiment of the invention, a method is
20 provided for determining whether a DNA substrate has a single nucleotide polymorphism (SNP). The method includes the steps of: contacting the DNA substrate with a modified DNA cleaving enzyme as described above; and determining from the cleavage product whether the DNA substrate has the SNP. It can further be
25 determined which nucleotide forms a SNP and the location of the SNP (see for example Figure 11)

 In a further embodiment of the invention, a method is provided of forming a shotgun cloning library, that includes the
30 steps of (a) incubating a modified DNA cleaving enzyme as described above with a DNA to form non-sequence specific cleavage fragments of the DNA that are ligatable; the ligatable DNA being capable of insertion into a vector for cloning in a host cell; and forming the shotgun cloning library.

5

In a further embodiment of the invention, a method for mapping nicks in a DNA is provided having the following steps: (a) incubating an enzyme according to the above with the DNA in a manganese-containing buffer; (b) permitting nicking to occur across from a pre-existing nick site to form double-stranded DNA with single strand overhangs; and (c) mapping the nicks in the DNA.

10

In a further embodiment of the invention, a method is provided for over-expressing T7 endonuclease 1, that includes selecting a host cell as above and over-expressing the T7 endonuclease 1.

15

BRIEF DESCRIPTION OF THE FIGURES

20

In the figure descriptions below, unless otherwise stated, "buffer" refers to standard buffer and Mn^{2+} or Mg^{2+} buffer refers to the standard buffer plus 2mM of the metal salt.

25

Figure 1 shows an SDS-PAGE analysis of expression and purification of unmodified MBP-T7 Endo I (ME) protein fusion.

30

Cultures of ER2566 containing pME, pME(PA/A) and pME(Δ PA) were grown at 37°C to mid-log phase, then at 30°C with 0.5 mM IPTG for 5 hours. The induced cells were harvested by centrifugation, suspended in standard buffer (20 mM Tris pH 7.6, 50 mM NaCl and 1 mM EDTA) and opened by sonication. After cell debris was removed by centrifugation, the crude cellular extracts were applied onto an amylose column. After the crude extracts flowed through, the column was washed with the buffer extensively.

35

- 5 The fusion protein was eluted with the buffer containing 10 mM maltose. SDS-PAGE was carried out on a 10-20% gradient gel. 30 μ l samples of the induced culture were used for each lane.

Lanes are labeled as follows: C, crude cellular extracts, F, the
10 flow through fraction and E, maltose elution fraction.

Figure 2 shows determination of enzyme activity by titration assay. 1 μ g of pUC(AT) in 20 μ l of either 2mM $MgCl_2$ or 2mM $MnCl_2$ in standard buffer was incubated with variable amounts of enzyme
15 at 37°C for 30 minutes. Adding DNA sample buffer on ice stopped the reactions. The samples were resolved on a 1.2% agarose gel with ethidium bromide.

Lane assignments in all panels are:

lane 1, no enzyme; lane 2, 5ng enzyme; lane 3, 2.5ng
20 enzyme; lane 4, 1.25ng enzyme; lane 5, 0.63ng enzyme;
lane 6, 0.32ng enzyme

Panels are:

Panel a1 shows ME(WT) in Mg^{2+} buffer

Panel a2 shows ME(WT) in Mn^{2+} buffer

25 Panel b1 shows ME(PA/A) in Mg^{2+} buffer

Panel b2 shows ME(PA/A) in Mn^{2+} buffer

Panel c1 shows ME(Δ PA) in Mg^{2+} buffer

Panel c2 shows ME(Δ PA) in Mn^{2+} buffer

30 Figure 3 shows cleavage of pUC(AT) by ME(Δ PA) in two step reactions.

In Panel a, pUC(AT) was incubated with the enzyme (DNA/ enzyme: 1 μ g/1.5ng) at 37°C in Mg^{2+} standard buffer. At variable

5 time points, an aliquot of sample was withdrawn and mixed with DNA sample buffer on ice. The collected samples were resolved on a 1.2% agarose gel with ethidium bromide.

In Panel b, the experiment was performed as in Panel A,
10 except the DNA/enzyme was 1 μ g /0.5 μ g.

Figure 4 shows the determination of non-specific nuclease activity by gel electrophoresis. 1 μ g of lambda DNA was incubated with variable amounts of enzyme in 20 μ l of either Mg²⁺ or Mn²⁺
15 buffer at 37°C for 30 minutes. The reaction was stopped by adding DNA sample buffer, then analyzed by agarose gel electrophoresis.

Lanes: lane 1, no enzyme added; lane 2, with 3ng of enzyme; lane 3, 16ng; 4, 80ng; 5, 400ng

In Panel a1, ME in Mg²⁺ buffer.

20 In Panel a2, ME in Mn²⁺ buffer.

In Panel b1, ME(PA/A) in Mg²⁺ buffer.

In Panel b2, ME (PA/A) in Mn²⁺ buffer.

In Panel c1, ME(Δ PA) in Mg²⁺ buffer.

In Panel c2, ME(Δ PA) in Mn²⁺ buffer.

25

Figure 5 shows the determination of nick site cleavage activity of the DNA cleaving enzyme.

Figure 5A shows pNBI plasmid having a nick site and a
30 restriction site. The plasmid was nicked by N.BstNB1 and then digested with BsaHI to yield a linearized double-stranded molecule containing a nick.

5 Figure 5B shows the results of different enzyme mixtures by gel electrophoresis.

 Lane 1, pNB I DNA;

 Lane 2, pNB I digested with N.BstNB I;

 Lane 3, with N.BstNB I, then with BsaH I;

10 Lane 4, with N.BstNB I and BsaH I, then with 10ng of ME in Mg^{2+} buffer at 37°C for 30 minutes;

 Lane 5, with BsaH I;

 Lane 6, with BsaH I, then with ME.

15 Figure 6 shows the determination of nick site cleavage in the presence of different metal ions. 1 μ g of the prepared substrate incubated with variable amounts of modified or unmodified enzyme at 37°C for 30 minutes in either Mg^{2+} or Mn^{2+} buffer. The digests were resolved on agarose gel.

20

 Lanes for all panels: Lane 1, no enzyme; Lane 2, with 2.5ng of enzyme; Lane 3, 5ng; Lane 4, 10ng; Lane 5, 20ng.

 In panel a1, the unmodified enzyme ME (WT) was used in Mg^{2+} buffer.

25 In panel a2, ME (WT) in Mn^{2+} buffer.

 In panel b1, ME(PA/A) in Mg^{2+} buffer.

 In panel b2, ME(PA/A) in Mn^{2+} buffer.

 In panel c1, ME(Δ PA) in Mg^{2+} buffer.

 In panel c2, ME(Δ PA) in Mn^{2+} buffer.

30

 Figure 7 shows the determination of the cleavage pattern at a DNA nick site by a modified DNA cleaving enzyme. Schematic representation of the method for analyzing the cleavage pattern at a nick site is provided.

5

Figure 8 shows the determination of DNA cleavage at single base mismatches by modified and unmodified DNA cleaving enzyme using hybrid mixtures as substrates. 0.5 μ g of each of hybrid mixture substrates was incubated with 20-30ng of enzyme in a 20 μ l reaction at 37°C for 30 minutes. The digests were resolved on an agarose gel.

Lanes for each panel: lane C, PCR product pcrAAT DNA or pcrACT as control; lane 1, hmAAT x ACT; lane 2, hmAAT x AGT; lane 3, hmAAT x ATT; lane 4, hmACT x AGT; lane 5 hmACT x ATT; and lane 6, hmAGT x ATT.

In panel a1, ME in Mg^{2+} buffer

In panel a2, ME in Mn^{2+} buffer.

In panel b1, ME(PA/A) in Mg^{2+} buffer.

In panel b2, ME(PA/A) in Mn^{2+} buffer.

In panel c1, ME(Δ PA) in Mg^{2+} buffer.

In panel c2, ME(Δ PA) in Mn^{2+} buffer.

Figure 9 shows how DNA substrates are generated for determination of enzyme activity on each individual single base mismatched DNA.

Panel A shows a schematic representation of preparation of an individual DNA heteroduplex with a definitive single-base mismatch. For making positive strand DNA, non-phosphorylated forward primer, pr-1 and phosphorylated reverse primer, P-pr-2 were used for PCR. The bottom strand, or the negative strand of the PCR product was removed by lambda exonuclease. The negative strand DNA was prepared by the same method as the positive one except

- 5 that phosphorylated forward primer P-pr-1 and non-phosphorylated reverse primer pr-2 were used.

Panel B is an example for generating a single-base mismatched heteroduplex by annealing two purified single stranded
10 DNAs

Lane 1, double stranded DNA produced by PCR, pcrAAT or pcrACT;

Lane 2, the positive (top) strand ssACT(+) isolated from pcrACT;

15 Lane 3, the negative (bottom) strand ssAAT(-) isolated from pcrAAT;

Lane 4, single-base mismatch (C/T) heteroduplex produced by annealing the purified positive strand with the negative one.

20 Figure 10 shows the determination of DNA cleavage by modified or unmodified DNA cleaving enzymes at individual single-base mismatches. 16 DNA duplexes were generated by annealing each one of the 4 isolated positive strands with each one of the 4 negative strands. Among them 4 were regular (perfect match) duplexes; 12 were heteroduplexes with single-base mismatches.
25 0.2-0.3 μg of the prepared duplex DNA were incubated with 10-15 ng of enzyme in 10 μl of either Mg^{2+} or Mn^{2+} buffer at 37°C for 20-30 minutes. The digests were resolved on agarose gels.

In panel a1 and a2, different duplexes were incubated with
30 ME in Mn^{2+} buffer.

In panel b1 and b2, they were incubated with PA(PA/A) in Mn^{2+} buffer.

5 Figure 11 shows the determination of the cleavage pattern by a modified DNA cleaving enzyme at base mismatch sites.

Panel a is a schematic representation of the experiment.

Panel b shows the experimental intermediate products resolved on a 1.2% low-melting agarose gel.

10 Lane 1, the mixed open plasmids after melt-anneal treatment;

Lane 2, after treatment with T4 ligase;

Lanes 3 and 4, after cleavage with ME(PA/A) in Mn^{2+} buffer followed by blunting the ends with T4 DNA polymerase. The arrow
15 symbol indicates the linear plasmid band that was excised and used for transformation after ligation.

Panel c shows a DNA sequence correlating to (a) and (b).

Figure 12 shows the three-dimensional structure of T7 Endo I
20 and the β -bridge. The homodimer is shown at left with one monomer displayed in light gray and the other in dark gray. At the right is a detail of the β -bridge showing the positions of the proline and alanine residues mutated in this study. Residues marked with
(b) refer to the second monomer (light gray).

25

Figure 13 shows the DNA and amino acid sequence for unmodified T7 Endo I.

Figure 14 are protein sequences from phage that have at
30 least 35% amino acid identity with the T7 Endo I sequence in Figure 13.

Figure 15 is a map of cleavage sites in lambda showing that nicking with modified T7 Endo I is random. Full length linear lambda

5 phage DNA was digested with the ME fusion enzyme containing the
PA to A sequence modification (ME[PA/A]) in the presence of 2mM
MnCl₂. DNA fragments from the reaction were purified by phenol
extraction and EtOH precipitation. After resuspension, the ends of
the DNA fragments were filled in using T4 DNA polymerase and
10 dNTPs (12°C or room temperature, 30 min). The blunt-ended
fragments were inserted into the EcoRV site of cloning vector
Litmus28 by adding aliquots of the ligation reaction to gel-purified
Litmus28 that was previously digested with EcoRV and
dephosphorylated with Calf Intestinal Phosphatase (CIP). The
15 ligation mixture was used to transform competent *E. coli* TB1 cells.
DNA was purified from ampicillin resistant transformants and the
inserts sequenced. Recovered sequence was examined to locate
vector-insert junctions and junctions between normally non-
contiguous fragments of lambda DNA. The endpoints of the
20 junctions were mapped to the complete lambda DNA sequence.
The distribution of cleavage sites on the lambda genome are shown
here using ME[PA/A] where this distribution is random.

25

DETAILED DESCRIPTION

Abbreviations: MBP, maltose binding protein, ME, maltose
binding protein - T7 endonuclease 1 conjugate, PCR, polymerase
30 chain reaction, PAGE, polyacrylamide gel electrophoresis, MCS,
multiple cloning site, dNTP, deoxyribonucleotide triphosphate, ATP,
adenosine triphosphate, SNP, single nucleotide polymorphism. The
term nuclease is used interchangeably with endonuclease.

5

"Cleavage" refers to a break in one or both of the strands of a duplex at opposite positions.

10 "Opposite positions" refers to sites on the top and bottom strands of the duplex DNA, where the sites are separated by less than 11 bases from each other.

The present embodiments apply to a class of enzymes characterized by a similar architecture to T7 Endo I and including T7
15 Endo 1 namely having two catalytic domains and a β -bridge (see for example, Figure 12). Enzymes in this class can be identified using the Conserved Domain Architecture Retrieval Tool (CDART) program of the National Center for Biotechnology Information (Geer, Lewis Y., et al. *Genome Research* 12:1619-1623 (2002)) or by other
20 predictive programs, based on searches employing the sequence of T7 Endo I (Figure 13).

Examples of enzymes identified in this manner include: T odd phages or related viruses including Enterobacteria phage T7,
25 NP_848275, endonuclease of Yersinia pestis phage phiA1122; NP_813757 putative endonuclease of Pseudomonas phage gh-1; NP_744417, phage endodeoxyribonuclease I of Pseudomonas putida KT2440; NP_523312, endonuclease of Bacteriophage T3; NP_064756, RP Endonuclease I of Roseophage SIO1; and
30 NP_052083, endonuclease of Bacteriophage phiYeO3-12. In addition other related phages such as SP6, bacteriophage phiKMV, Enterobacteria phage K1-5, Vibriophage VpV262, BA14, BA127 and BA156 may encode similar enzymes.

5 Examples of enzymes having at least 35% amino acid identity with T7 Endo I are listed in Figure 14.

 The reciprocal stereo-geometric positions of the two catalytic centers in an enzyme of the above class can be changed by
10 introducing mutations into the bridge without changing the catalytic centers per se, using genetic or biochemical means. Consequently, as exemplified herein, mutations were introduced into the β -bridge site without interfering with the folding and function of the catalytic domain (Example 1). Embodiments of the invention are not
15 intended to be limited by (a) the exemplified methods for introducing mutations into sites in the β -bridge or (b) any particular location of the mutation within the bridge. The term "mutation" is intended to mean deletion, substitution or addition of one or more amino acids in the enzyme.

20

 Mutations in the bridge portion of the enzyme resulted in shifting the enzyme activity profile to different substrates and in alteration of the reaction kinetics and distribution of the final products. Additionally, mutations in the bridge portion result in
25 altered enzyme activity profiles in different salt solutions. A practical consequence of the observed changes in properties of a modified enzyme compared with unmodified enzyme is the reduced toxicity of the enzyme activity *in vivo* while enzyme activity is preserved or enhanced *in vitro*. Consequently, the modified
30 enzyme can be cloned and over-expressed in host cells because of its reduced toxicity to the host cell. This makes it possible to produce amounts of enzyme suitable for its practical use as a reagent for molecular biology applications.

5 The change in the activity profile of the modified enzyme as a result of one or more mutations was achieved when (i) one or more mutations were introduced in the β -bridge segment that connects the two well-separated catalytic domains, and/or (ii) the reaction conditions were manipulated. For example, the metal ion
10 composition of the reaction buffer was selected to provide an altered activity profile for the enzyme. In the Examples, magnesium or manganese ions were included in the reaction buffer. These buffers may also be used together to further modify the enzyme activity of the mutants. For example, if manganese ions are
15 added to the reaction mixture containing magnesium ions, the non-sequence specific nicking can be reduced. The use of metal ions other than magnesium or manganese, either alone or together with magnesium or magnesium at various concentrations in the reaction buffer is not precluded.

20

 In embodiments of the invention, computational analysis of the T7 Endo I structure and in particular the β -bridge revealed that the bridge constitutes part of the dimer interface, and that residues are bound to each other by hydrogen bonds except for the two
25 residues, P46 and A47, at the bridge center (Figure 12). The two residues are located at the geometrical center of the protein, rendering it flexible.

 Genetically changing the distance between the two catalytic
30 centers was achieved by introducing mutations into the β -bridge site of the protein. For example, for T7 Endo I, PA/A, PA/AA, PA/PGA, PA/PAPA, Δ PA, PA/K, PA/G, PA/D and PA/P mutants were made with altered enzyme activity which was established in reaction buffers containing magnesium or manganese ions using a variety of

5 DNA substrates including mismatch duplex DNA, duplex DNA containing nicks, un-nicked duplex DNA and DNA having cruciform structures. The conformational flexibility of a protein having a structure similar to T7 Endo I contributes to the broad substrate specificity of the enzyme. Surprisingly, nucleotide substitutions in
10 the β -bridge that were of variable length (single amino acid, dipeptide, tripeptide or tetrapeptide), negatively charged, positively charged, or neutral in charge, flexible in structure or inflexible in structure revealed altered enzyme activities of the type described herein.

15

Approximately twenty-four β -bridge site mutants were generated. The proteins were purified and characterized (see for example, Example 1) All the mutations in some way altered the activity profile of the enzyme. Two active deletion mutants,
20 ME(PA/A) and ME(Δ PA), were characterized extensively (see Example 1). As expected, the two catalytic domains in the mutants were folded correctly and were fully active, since both could resolve cruciform DNA with an efficiency similar to that of the wild type protein. For resolving cruciform structures, the wild type enzyme
25 required Mg^{2+} for its activity. The activity was reduced by 20 fold or more if Mg^{2+} was replaced with Mn^{2+} . For the two mutants, the metal ion requirement was relaxed. The mutant proteins were almost equally active in either Mg^{2+} or Mn^{2+} buffer. However, the kinetics and the final products with ME(Δ PA) were changed. This
30 mutant could no longer simultaneously cleave both strands across the cruciform structure on plasmids. Instead, it nicked one strand first at the site in a rapid reaction, resulting in relaxation of the cruciform, then cleaved the other strand at the generated nick in a separate slow reaction.

5

For cleavage of DNA at nicks or base mismatches and for the non-sequence specific nuclease activity, the requirements for metal ions among the enzyme and its mutants were shifted again. This time the wild type ME remained about equally active in either Mg^{2+} or Mn^{2+} buffer. For the mutants, however, these activities, were significantly reduced in Mg^{2+} buffer. Mn^{2+} became the required divalent metal ion. In Mn^{2+} buffer, ME(PA/A) could recognize and cleave DNA at nick sites and at certain types of single-base mismatch sites even more efficiently than its wild type counterpart.

15

One of ordinary skill in the art will appreciate that a duplex DNA can be described as having a top strand and a bottom strand and that a mismatch occurs when a nucleotide at a position on the top strand is not complementary to the nucleotide at the same position on the bottom strand. Cleavage with the DNA cleaving enzyme can cause at least one of (a) a break at a 3' flanking site of the mismatched nucleotide on the top and bottom strands or (b) a break at the 5' flanking site of the mismatch on the top and bottom strands where the flanking site is preferably less than about 11 nucleotides from the mismatch.

25

Mutants of enzymes in which amino acids in the bridge have been altered or deleted can be significantly less toxic than the wild type enzymes *in vivo*. Mutants such as ME (PA/A) appear to have significantly reduced non-specific nuclease activity in the presence of Mg^{2+} (present in the host cell in millimolar concentrations). This makes it possible to over-express a mutant T7 endonuclease I such as ME(PA/A). Moreover, the mutant has enhanced resolvase activity

30

- 5 in Mn^{2+} (not present in the host cells) compared with the wild-type enzyme.

Enzymes belonging to the T7 endonuclease class (including proteins with at least 35% identity in amino acid sequence), in
10 which one or more mutations have been introduced into the protein outside the catalytic domains and preferably within the bridge between the catalytic domains, have advantageous properties. Examples of advantageous properties as exemplified by T7 Endo I are:

15

1. The unmodified enzyme is toxic to cells while the modified enzyme is not thereby permitting cells to overexpress the mutant enzyme (Example 1)

20

2. Cleavage of cruciform structures. The unmodified enzyme ME(WT) had 20 fold reduction of activity in manganese buffer compared with its activity in magnesium buffer. In contrast, modified enzymes ME (PA/A) and ME (Δ PA) showed similar activity in the different buffers. However, ME (Δ PA) generated nicked DNA
25 or linear DNA in equal amounts whereas ME(PA/A) produced double strand breaks (Example 2). This altered activity permits resolution of branched DNA.

30

3. The unmodified and the modified DNA cleaving enzymes both can find a nick in double-stranded DNA and cleave against the nick resulting in non-sequence specific cleavage of double-stranded DNA into fragments with 3 or 4 base overhangs (Example 2). However, the modified enzyme has greater cleavage activity in

5 manganese buffer than in magnesium containing buffer. This property has utility in nick site mapping.

4. Both modified and unmodified enzymes have non-specific nuclease activity but the modified enzyme has less nuclease activity
10 in magnesium containing buffer than the wild-type enzyme. The modified enzyme has greater nuclease activity of an amount comparable to the unmodified enzyme in manganese-containing buffers. Non-specific nuclease cleavage can provide random ligatable DNA sequences suitable for shotgun or other types of
15 cloning. An advantage of this approach over the use of, for example, DNase I is that it is possible to control the size of the fragments and to avoid shredding the DNA of interest into tiny pieces.

20 5. Both modified and unmodified enzymes recognize single base mismatches. However, the modified enzyme has greater activity and recognizes a wider range of substrates than the unmodified enzyme in manganese buffer. For analysis of single nucleotide polymorphisms (SNPS), the target DNA is denatured and
25 hybridized to related DNA, which lack the polymorphisms. This results in single base mismatches of which there are 8 possibilities: A to A, A to G, A to C, C to T, C to C, G to G, G to T and T to T. These can all be recognized by the modified enzyme (Example 2). These modified enzymes provide a useful tool for SNP detection
30 analysis utilizing mismatch detection.

A DNA cleaving enzyme having at least 35% amino acid identity with T7EndoI and having two catalytic centers separated by a β -bridge, where the β -bridge contains a mutation that results in

5 altered enzyme activity as described herein can be provided in a kit with a suitable buffer, for example, containing manganese ions or magnesium ions as appropriate for the desired function according to above. The kit may additionally contain a DNA size ladder. Instructions may further be included in the kit. Alternatively, the kit
10 may include a plasmid encoding the endonuclease or indeed a transformed host cell preparation containing a recombinant vector expressing the mutated nicking endonuclease.

All references cited herein are incorporated by reference. In
15 addition, U.S. provisional application serial number 60/524,123 filed November 21, 2003 and Guan et al. *Biochemistry* 43:4313-4322 (2004) are herein incorporated by reference.

20

EXAMPLES

Example 1: Generation of Mutations at the Bridge Site and Over-expression of the Modified Enzymes

25

Materials and Methods

Restriction enzymes, nicking enzyme N.BstNB I, DNA
polymerases, T4 ligase, T4 DNA kinase, β -agarase, λ Exonuclease,
the maltose-binding protein (MBP) protein fusion expression and
30 purification system including plasmid pMAL-c2x, the host *E. coli*
strains TB1 and ER2566, Factor Xa protease, the cruciform
structure-containing plasmid pUC(AT) and plasmid LITMUS28 were
obtained from New England Biolabs Inc. (New England Biolabs, Inc.,
Beverly, MA). Synthetic oligonucleotides were synthesized using

- 5 standard techniques. T7 phage DNA encoding T7 Endo I has a sequence corresponding to SEQ ID NO:1.

Method for Recombinant DNA and Mutagenesis

- 10 DNA manipulation and site-directed mutagenesis (Kunkel, *Proc. Natl. Acad. Sci. U S A.* 82(2):488-92 (1985 Jan)) or by PCR were carried out as described in Molecular Cloning by Sambrook et al. (Sambrook, J. and Russell, D.W. (2001) Molecular Cloning: a Laboratory Manual, pub. Cold Spring Harbor Laboratory, Cold
15 Spring Harbor, NY (2001)).

Cloning T7 EndoI mutants

- For cloning T7 Endo I from phage chromosomal DNA by PCR,
20 two primers were used:

oligo-1: CCCGAATTCATGGCAGGTTACGGCGCT (SEQ ID NO:2;
and

- oligo-2: CCCCCAAGCTTATTTCTTTCCTCCTTT (SEQ ID NO:3)
25

- The PCR product was treated with restriction enzymes, and cloned into the Hind III-EcoR I site of plasmid pUC19, resulting in plasmid pEndo I. For construction of pEndo($\Delta\beta 2$) (where $\Delta\beta 2$ is deletion of the second β sheet) and pME($\Delta\beta 2$) from pEndo I, two
30 sequential PCRs (two-step PCR) were performed. In the first step pEndo I was used as template.

Oligo-1 and oligo-4
(TGGAAGTAAGAAGTCTGGCCACTCTTCATA) (SEQ ID NO:4) were

- 5 used as primers for obtaining the 5' part of the gene; oligo-3 (TTCGAGTATGAAGAGTGGCCAGACTTCTTA) (SEQ ID NO:5) and oligo-2 were used for the 3' part of the gene.

10 In the second step, a 1 to 1 mixture of the purified 5' and 3' products from the first step was used as template; oligo-1 and -2 were used as primers. For each step, 10 to 15 cycles (95°C, 0.5 min; 45°C, 0.5 min; 72°C, 0.5 min) of PCR were performed. The final product was cloned into pUC19 and pMAL-c2x resulting in pEndo($\Delta\beta$) and pME($\Delta\beta$ 2), respectively. A recognition sequence for
15 restriction enzyme Msc I was generated at the deletion site. For construction of pME(PA/A) from pME($\Delta\beta$ 2), two DNA oligonucleotides,

oligo-5: AAAGTGCCTTATGTAATTGCGAGCAATCACACTTACACT
20 (SEQ ID NO:7)

oligo-6: AGTGTAAGTGTGATTGCACGCAATTACATAAGGCACTTT
(SEQ ID NO:8)

were annealed, then inserted into the Msc I site of pME($\Delta\beta$ 2).
25 pME(Δ PA) was constructed as pME(PA/A), except for using two different oligonucleotides,

oligo-7: AAAGTGCCTTATGTAATTAGCAATCACACTTACACT (SEQ
ID NO:9) and oligo-8:
30 AGTGTAAGTGTGATTGCTAATTACATAAGGCACTTT (SEQ ID
NO:10).

5 Preparation of Substrates for Heteroduplex Analysis

For construction of pAAT, pACT, pAGT and pATT from pEndo($\Delta\beta 2$), two oligonucleotide mixtures were:

- 10 oligomix-9: AAAGTGCCTTATGTAAATTCCTAATCACACTTACACT
(SEQ ID NO:11); and oligomix-10:
AGTGTAAGTGTGATTANTGGGAATTTACATAAGGCACTTT (SEQ ID
NO:12) were annealed, then inserted in the Msc I site of
pEndo($\Delta\beta 2$). The desired individual clones were verified by DNA
15 sequencing.

DNA sequencing was performed on Applied Biosystems, Inc.'s automated DNA sequencers (3100) using Big Dye labeled dye-terminator chemistry (Applied Biosystems, Inc., Foster City, CA).

- 20 The cloned DNA and the DNA used as templates for preparing substrates by PCR were verified by sequencing.

Preparation of Heteroduplex Substrates

- 25 For preparation of hybrid substrates by melt-anneal treatment, two purified PCR products described above were mixed in annealing buffer (20mM Tris. 7.6, 50 mM NaCl) at a 1:1 ratio. The mixture was incubated at 95°C for 5 minutes followed by incubating at 65°C for 1 hour, then gradually cooled down to room
30 temperature. For preparation of heteroduplexes by annealing purified single-strand DNA, a positive strand and a negative strand were mixed at 1 to 1 ratio in annealing buffer, then subjected to melt-anneal treatment.

5 Gene Expression and Protein Purification

Expression and purification of gene products using the MBP fusion and expression system were carried out as described previously (Sambrook, J. and Russell, D.W. (2001) Molecular Cloning: a Laboratory Manual, pub. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (2001); Riggs, P. Current Protocol in Molecular Biology, ed. Ausubel, F.A. et al. 16.6.1-16.6.14, pub. Green Associates/Wiley Interscience, New York (1994)).

10 Modifications to the standard protocol and preparation of non-fusion enzyme were described previously (Guan, C., et al. *J. Biol. Chem.* 19:1732-1737 (1996)). The purified enzyme, either the native enzyme or its MBP fusion form, was stored in 50% glycerol at -20°C. Protein Analysis-Protein concentration was determined by the Bio-Rad Protein Assay using bovine serum albumin as a

15 standard. Molecular weight and purity determinations were carried out by SDS-PAGE analysis and MALDI-ToF mass spectrometry using Applied Biosystems Inc.'s Voyager DE (Applied Biosystems Inc., Foster City, CA). N-terminal protein sequence analysis was performed on a Procise 494 Protein/Peptide Sequencer (Applied

20 Biosystems Inc., Foster City, CA).

25 Biosystems Inc., Foster City, CA).

To assay resolvase activity, 1 µg of the cruciform-containing plasmid pUC(AT) in 20 µl of buffer (20 mM Tris, pH 7.6, 50 mM NaCl, 2 mM DDT) with 2 mM MgCl₂ or 2 mM MnCl₂ was incubated

30 with variable amounts of purified enzyme at 37°C for 30 minutes. The digests were then analyzed by agarose gel-electrophoresis. Quantitative analysis of DNA or protein bands on the gel was performed on a Bio-Rad Phosphoimage autodensitometer.

5 Results: Preparation and Overexpression of T7 Endonuclease
 Mutants

 The T7 Endo I gene was obtained by PCR using genomic DNA isolated from T7 phage as a template. The gene was cloned as a
10 480-base pair (bp) EcoRI-HindIII fragment on plasmid pUC19, named pEndo I, with the orientation of the cloned gene opposed to the lac promoter on the vector. This EcoRI-HindIII fragment should be transferable from pEndo I into the EcoRI-HindIII site of pMAL-c2x plasmid to form an MBP-Endo I in-frame gene fusion. The first
15 attempt to construct the MBP-Endo I fusion in the pMAL system was unsuccessful. The ligation mixture yielded no colonies upon transformation of the host cells, even though multiple copies of the lacIq gene were present in the host. Anticipating that this failure might result from toxic expression even in un-induced cells, a
20 variant pMal vector (pMalC2x-NEB) was constructed with the more-tightly regulated T7 promoter instead (Studier et al. *Meth. Enzymol.* 185:60-89 (1990)). Using this vector, the construct was obtained. In a T7-promoter expression host such as ER2566 or TD-1, 2 to 5 μ g of ME could be purified routinely from 1 liter of induced culture.

25

 To generate mutations in the β -bridge while minimizing PCR errors, we constructed an inactive recipient ME fusion vector with the region of interest deleted, to which synthetic oligonucleotides could be added back. Briefly, the DNA encoding the second β sheet
30 (β 2), from residue 40 to 53, was removed from pEndo I by two-step PCR, and replaced by an MscI site. The resulting plasmid was named pEndo($\Delta\beta$ 2). The mutated gene Endo($\Delta\beta$ 2) was transferred as an EcoR I-Hind III fragment into the EcoR I-Hind III site of pMAL-c2x to produce plasmid pME($\Delta\beta$ 2). This plasmid was stable in

5 the *E. coli* host TB1 or ER2566. Other mutants were generated by inserting short synthetic double-stranded DNA into the MscI site of pME($\Delta\beta 2$). The MBP-Endo I mutants were named as follows: ME(Δ PA), where the two residues P46 and A47 located at the β -bridge center were removed; ME(PA/A) where the dipeptide PA was
10 replaced by the single residue A. The dipeptide PA was also replaced by other single amino acids, dipeptides, tripeptides or tetrapeptides to generate different variants such as ME(PA/G), ME(PA/AA), ME(PA/PGA) and ME(PA/PAPA) and so on. In mutation ME(Δ bdg), all 6 residues (44-49) that form the bridge were removed.

15

The severe cellular toxicity of T7 EndoI was significantly reduced in many of these mutants, since the mutant genes could be maintained in pMALc-2-x vectors without causing severe host growth defects. In contrast to the wild-type ME, large amounts of
20 MBP-fused mutant proteins were produced after induction. 30-50 mg/L of fusion protein was routinely obtained following a single step of amylose affinity chromatography (Figure 1).

The purified wild-type ME fusion was as active as the enzyme
25 from which the MBP part had been removed using Factor Xa cleavage, judged by activity titration using pUC(AT) (see Materials and Methods above and discussion below), and taking the molecular weight of fusion protein into account. Since the MBP-Endo I fusion was fully active, the experiments in this study were carried out
30 using the purified fusion proteins.

5 **Example 2: Characterization of two deletion mutants**
ME(PA/A) and ME(Δ PA).

1. Cleavage of Cruciform Structure DNA

10 A cruciform structure stabilized on a negatively supercoiled plasmid is structurally similar to a four-way junction of DNA. T7 Endo I resolves both DNA structures with high efficiency (Parkinson, M.J. and Lilley, D.M.J. *J. Mol. Biol.* 270:169-178 (1997)). To compare the enzymatic activities among different T7 Endo I
15 mutants, we defined specific activity using cleavage of a cruciform-containing-plasmid, pUC(AT) as a substrate. One unit of activity was defined as the amount enzyme that was needed to convert 1 μ g of supercoiled pUC(AT) to either the linear form (with both strands cut) or the nicked form (with a single strand cut), in 20 ml
20 of reaction at 37°C in 30 minutes. We measured the specific activity by enzyme titration assay. The results are presented in Figure 2.

In Mg^{2+} buffer, the specific activities of ME and ME(PA/A) were about the same, 600 to 800 U/mg (Figure 2, Panels a1 and
25 B1). When Mg^{2+} buffer was replaced with Mn^{2+} buffer, the activity of ME was reduced by about twenty-fold, to only about 40 U/mg (Figure 2, panel a2). In contrast, the activity for ME(PA/A) remained about the same (600 to 800 U/mg) in either Mg^{2+} or Mn^{2+} buffer (Figure 2, panels b1 and b2). The final products with either ME or
30 ME(PA/A) were basically the same, linearized plasmid. Singly-nicked intermediates were not observed, suggesting that cleavage of both strands is concerted. Thus, the deletion of proline relieved the Mn^{2+} inhibition without altering the reaction progress.

5 In contrast, for ME(Δ PA), a change in reaction progress and in
ion-dependence was observed. The overall activity of this enzyme
was slightly lower, about 400 U/mg, and remained about the same
in either Mg^{2+} or Mn^{2+} buffer (Figure 2, Panels c1 and c2). However,
the products were dramatically different from those produced by
10 wild-type and ME(PA/A): in Mg^{2+} buffer most ($\geq 90\%$) of the final
products were nicked plasmids (single cleavage) and less than 10%
were linear products (double cleavage; Figure 2, Panel c1). For this
enzyme but not the others, Mn^{2+} affected the identity of the
products, increasing the percentage of doubly-cleaved products to
15 50% (Figure 2, Panel c2).

 We examined the progress of the reaction both at low and at
high enzyme: substrate ratio in Mg^{2+} buffer (Figure 3). Although
ME(Δ PA) could eventually convert the nicked plasmid into linear
20 form after a prolonged incubation, or when high concentrations of
enzyme were used (Figure 3, Panel b), at low enzyme concentration
the rate of second-strand cleavage was more than 100 times slower
than that of the first one. Reaction conditions in substrate excess
(with low concentrations of enzyme, as in Figure 3, Panel a) require
25 each enzyme molecule to carry out multiple turnovers to complete
the first strand cleavage on all substrate molecules. Thus, first and
second strand cleavage appear to be two separate reactions for this
enzyme.

30 Restriction digestion analysis on the final products of the
resolvase reaction confirmed that ME(PA/A) and ME(Δ PA) cleaved
pUC(AT) at the cruciform structure site as did ME. Further
shortening of the bridge to yield ME(Δ bdg) resulted in very low

- 5 nicking activity (5-10 U/ μ g) without any double-stranded cleavage. This residual activity was still specific for the cruciform structure.

These results showed that the two catalytic domains in the active mutants were folded correctly and fully active. We also infer
10 that each domain functions independently as a nicking endonuclease. It was necessary to maintain a proper geometrical relation between the two catalytic centers for the enzyme to effectively resolve cruciform DNA. Genetically changing the relative position of the two centers resulted in alteration of the enzyme
15 efficiency, the metal ion preference, the reaction kinetics and the distribution of final products.

2. Non-Specific Endonuclease Activity

20 Non-specific endonuclease activity of T7 Endo I was observed when high concentrations of enzyme were used. To compare the non-specific nuclease activity among T7 Endo I variants, 1 μ g of lambda phage DNA was incubated with variable amounts of enzyme in a 20ml reaction at 37°C for 30 minutes. The digests were
25 subjected to gel electrophoresis. The results are presented in Figure 4.

For the wild type enzyme ME, the non-specific nuclease activity was about the same in either Mg^{2+} or Mn^{2+} buffer (Figure 4,
30 Panel a1 and Panel a2). This contrasts with the 20-fold ion effect with the specific substrate (see above). Mutations also selectively affected the nonspecific nuclease activity: both mutants ME(PA/A) and ME(Δ PA), showed 20 and 50 times lower non-specific nuclease activity in Mg^{2+} buffer than did ME (Figure 4, Panels b1 and c1),

5 whereas on the specific substrate there was little reduction in
specific activity. Mn^{2+} spared the non-specific nuclease activity of
both mutants: nonspecific degradation was achieved at similar or
lower enzyme concentrations as observed with ME (Figure 4, Panels
b2 and b2).

10

Adding Mg^{2+} to the reaction mixture containing Mn^{2+}
competitively inhibited the elevated non-specific nuclease activity of
ME(PA/A) and ME(Δ PA). Addition of 10 to 20 mM of Mg^{2+} to the
2mM Mn^{2+} buffer eliminated more than 90% of the non-specific
15 nuclease activity of these enzymes.

20

Shotgun cloning and sequence analysis of lambda DNA
fragments produced by digestion with ME or ME(PA/A) indicated
that the enzyme cleaved linear DNA randomly (Figure 15).

3. Cleavage of Nicked DNA Duplex

We showed above that ME(Δ PA) cleaves DNA at a nick site,
eventually converting the nicked intermediate of pUC(AT) to the
25 linear form (Figure 3, Panel b). To determine whether T7 Endo I
could cleave at any nick or only during the progress of a resolution
reaction, we made a linear nicked substrate with no cruciform
character. Plasmid pNB I (New England Biolabs, Inc., Beverly,
MA), which contains a single site for the site-specific nicking
30 enzyme N.BstNBI, was digested with the nicking enzyme first, then
digested with BsaHI. The resulting product was a 2.5-kb linear DNA
molecule with a nick about 600 bp from one end of the molecule.
The same protocol without N.BstNBI nicking yielded an un-nicked
control substrate. These substrates were then incubated with ME in

5 Mg^{2+} buffer and fragments were resolved on an agarose gel. The nicked substrate yielded the expected 0.6-kb and 1.9-kb DNA fragments in addition to the original 2.5-kb DNA (Figure 5), while the control yielded only the 2.5-kb substrate DNA. This indicates that T7 Endo I recognizes the presence of a nick in the duplex and
10 cleaves the molecule at or near that site.

Cleavage opposite or approximately opposite a nick is a novel activity, so we investigated the properties of the mutants described above on this new substrate, and again compared the ability of
15 Mg^{2+} or Mn^{2+} to support the reaction. The results are presented in Figure 6.

In contrast to cruciform resolution, where Mn^{2+} inhibited the wild-type enzyme, the wild-type ME showed equivalent activity at
20 the nick in either Mg^{2+} or Mn^{2+} buffer (Figure 6, Panels a1 and a2). This suggests that it is the first-strand cleavage of the cruciform that is inhibited by Mn^{2+} for this enzyme. For ME(Δ PA), the activity in Mg^{2+} buffer was undetectable (Figure 6, Panel c1), while in Mn^{2+} the activity was significantly increased (Figure 6, Panel c2). This
25 behavior is compatible with its activity on the cruciform substrate, in that Mg^{2+} supported activity on the cruciform resulted primarily in nicked products, while Mn^{2+} resulted in double strand cleavage. This suggests that the enzyme was unable to efficiently complete the resolution reaction by cleaving opposite or approximately
30 opposite to the first nick in Mg^{2+} , but that Mn^{2+} supported cleavage opposite or approximately opposite the nick, as seen here.

In the presence of Mn^{2+} , ME(PA/A) was a more active enzyme than its wild-type counterpart for nick site cleavage. Indeed,

5 additional products are obtained, not seen with the wild-type where the products corresponded to the increased general non-specific activity observed with this enzyme above. In the present experiment, the amount of enzyme is much less (20 ng) than that used to obtain extensive non-specific activity (400 ng).

10

Figure 7 illustrates our approach to determining the location of cleavage of this nick site activity. Briefly, plasmid pNB1 was first treated with nicking enzyme N.BstNB I, then cleaved with ME or ME(PA/A) in Mn^{2+} buffer. The linear plasmids were identified and
15 purified from the agarose gel. After the ends were polished to create flush ends with T4 DNA polymerase plus dNTP, a 300 bp flush-ended Stu I-SnaB I fragment isolated from plasmid Litmus 28 and containing the multiple cloning site and primer binding sites for sequencing was ligated to these products. The recombinant
20 plasmids were isolated from individual transformants and sequenced using the two primers selected from oligonucleotides S12050 and S12051 (NEB, Beverly, MA), reading outward from the center of the MCS inserts. The results showed that the endonuclease introduced a single cut, in the continuous strand,
25 displaced 5' from the nick by 3 or 4 bp. These two positions were found with about an equal frequency (data not shown). The sequencing data also showed that under the conditions of the experiment, the enzyme did not cleave the short single-stranded overhangs at sticky ends generated during the reaction, since the
30 duplication created by the fill-in reaction was intact.

5 4. Cleavage of DNA Heteroduplex with Single-Base Mismatches

To test the ability of T7 Endo I and its mutants to cleave DNA at single-base mismatch sites, we made use of mutated pEndo I genes created during the mutagenesis project. Four individual pEndo I
10 derivatives, named pAAT, pACT, pAGT and pATT were constructed (see experimental procedures). The only difference among them was that there was a single base variation at the same position of the gene. Using the 4 individual clones as templates 4 PCR
15 products, about 480 bp long, named pcrAAT, pcrACT, pcrAGT and pcrATT, were obtained. Single-base mismatch heteroduplex molecules could be obtained by mixing any two of the 4 PCR products followed by melt-anneal treatment (Babon, J.J., et al. *Molecular Biotechnology*, 23:73-81 (2003)). 6 hybrid mixture
20 substrates, named hmAATxACT, hmAATxAGT, hmAATxATT, hmACTxAGT, hmACTxATT and hmAGTxATT, were prepared. Each hybrid mixture was expected to contain two different heteroduplexes besides the two original homoduplexes. For example, hmAATxACT contained both A/G and C/T single-base mismatch heteroduplexes besides the original pcrAAT and pcrACT.
25 The mismatch site was about 150 bp away from one end of the molecule. The maximum content of heteroduplexes in an individual hybrid mixture was 50%. Two different hybrid mixtures may contain the same kind of base mismatch, but the sequences flanking the mismatch site are different.

30

The 6 hybrid mixtures were incubated with ME, ME(PA/A) and ME(Δ PA), respectively, in either Mg^{2+} or Mn^{2+} buffer. The digests were subjected to agarose gel electrophoresis (Figure 8). The results showed that in all cases, ME digested the heteroduplex to

5 create 150-bp and 330-bp from the original 480-bp one. In the control directly using pcrAAT, ACT, AGT or ATT as substrates, only the original 480-bp DNA band could be identified (Figure 8, panels a1 and a2). These indicated that ME cut at least one of the two heteroduplexes present in each of 6 hybrid mixtures. The cleavage
10 efficiency on these substrates with ME was about the same in either Mg^{2+} or Mn^{2+} buffer. However, in Mg^{2+} buffer the cleavage with ME(PA/A) or ME(Δ PA) was almost undetectable (Figure 8, Panels b1 and c1). The cleavage efficiency for ME(PA/A) or ME(Δ PA) was significantly increased by replacing Mg^{2+} with Mn^{2+} in the reaction
15 (Figure 8, Panels b2 and c2). In the presence of Mn^{2+} , ME(PA/A) was at least as active as ME was for cleavage of single-base mismatched DNA in hybrid mixture substrates. ME(Δ PA) was a less active enzyme, even in Mn^{2+} buffer.

20 In order to assess the cleavage efficiency of the enzyme with each kind of single-base mismatch, we prepared heteroduplex substrates, each of which contained only one of the possible mismatches (Figure 9, Panel A). Briefly, 8 single-strand DNA molecules, about 480 base long, named ssAAT(+), ssAAT(-),
25 ssACT(+), ssACT(-) and so on were generated, using lambda exonuclease to selectively digest one strand of the relevant PCR product. Annealing each one of the positive strands with each one of the negative strands could produce 16 duplex DNA molecules, 12 of them single-base mismatch heteroduplexes and 4 homoduplex
30 molecules (Figure 9, Panel B). The base mismatch site was about 150 bp away from one of the ends of the molecules. Some heteroduplexes may contain the same kind of mismatches, but the sequences around the mismatches are different.

5 The 16 DNA duplexes prepared above were incubated with ME and ME(PA/A) in Mn^{2+} buffer respectively. The digests were resolved on agarose gels (Figure 10). The results showed that the 4 regular DNA duplex molecules generated by annealing two corresponding positive and negative strands were not specifically
10 cleaved by either ME (Figure 10, Panel A) or ME(PA/A) (Figure 10, Panel B). Both enzymes could effectively cleave heteroduplexes at the single-base mismatch sites where at least one of the mismatched bases was cytosine. Neither enzyme cleaved DNA at a G/G mismatch site very efficiently. The most profound difference
15 between the two enzymes was that ME(PA/A) had much higher activity at A/A and T/T mismatches than did ME. In general, ME(PA/A) in the presence of Mn^{2+} was a more efficient enzyme than its wild type counterpart when cleaving DNA at single-base mismatch sites. To determine the cutting pattern by the enzyme at
20 single-base mismatch sites, we performed an experiment as illustrated in Figure 11, Panel a. Briefly, plasmids pAAT, pACT, pAGT and pATT were linearized by digestion with HindIII. The four linear plasmids were mixed and subjected to melting-anneal treatment. The annealed plasmid mixture was treated with T4 ligase plus ATP
25 to produce relaxed circular molecules. The majority (up to 75%) of the recircularized plasmids should contain a single-base mismatch site that is sensitive to cleavage by T7 Endo I. The ligation mixture was digested with ME(PA/A) in Mn^{2+} buffer followed by treatment with T4 DNA polymerase plus dNTP to blunt the ends, then
30 subjected to agarose electrophoresis. The DNA band representing the full-length linear plasmids was isolated from the gel (Figure 11, Panel b). The purified linears were treated with T4 ligase and transformed into an *E. coli* host TB1. Plasmids were isolated from individual transformants sequence across the ligation junction using

- 5 a primer located 150 bp away from the original mismatch site. The results showed that ME(PA/A) cleaved both strands of DNA 5' to, and 1 to 3 bp away from, the mismatched base, generating 3 to 7 (many of them 4 to 6) bp long sticky ends, as judged by the length of the duplication generated at the mismatch site.